

Asthma Action Plan

ENGLISH/VIETNAMESE

PROVIDER INSTRUCTIONS

At initial presentation, determine the level of asthma severity

- Level of severity is determined by both impairment and risk and is assigned to the most severe category in which any feature occurs.



At subsequent visits, assess control to adjust therapy

- Level of control is determined by both impairment and risk and is assigned to the most severe category in which any feature occurs.
- Address adherence to medication, inhaler technique, and environmental control measures.
- Sample patient self-assessment tools for asthma control can be found at <http://www.asthmacontrol.com/index.html>
<http://www.asthmacontrolcheck.com>



Stepwise approach for managing asthma:

- Therapy is increased (stepped up) if necessary and decreased (stepped down) when possible as determined by the level of asthma severity or asthma control.

Asthma severity and asthma control include the domains of current impairment and future risk.

Impairment: frequency and intensity of symptoms and functional limitations the patient is currently experiencing or has recently experienced.

Risk: the likelihood of either asthma exacerbations, progressive decline in lung function (or, for children, reduced lung growth), or risk of adverse effects from medication.

ASTHMA MANAGEMENT RECOMMENDATIONS:

- Ensure that patient/family receive education about asthma and how to use spacers and other medication delivery devices.
- Assess asthma control at every visit by self-administered standardized test or verbal history.
- Perform spirometry at baseline and at least every 1 to 2 years for patients \geq 5 years of age.
- Update or review the Asthma Action Plan every 6 to 12 months.
- Perform skin or blood allergy tests for all patients with persistent asthma.
- Encourage patient/family to continue follow-up with their clinician every 1 to 6 months even if asthma is well controlled.
- Refer patient to a specialist if:
 - there are difficulties achieving or maintaining control
OR
 - step 4 care or higher is required (step 3 care or higher for children 0-4 years of age) OR
 - immunotherapy or omalizumab is considered OR
 - additional testing is indicated OR
 - if the patient required 2 bursts of oral systemic corticosteroids in the past year or a hospitalization.

HOW TO USE THE ASTHMA ACTION PLAN:

Top copy (for chart):

- File this copy in the patient's medical chart.

Middle copy (for patient):

- Enter specific medication information and review the instructions with the patient and/or family.
- Educate patient and/or family about factors that make asthma worse and the remediation steps on the back of this form.
- Complete and sign the bottom of the form and give this copy of the form to the patient.

Bottom copy (for school, childcare, work, etc):

- Educate the parent/guardian on the need for their signature on the back of the form in order to authorize student self-carry and self-administration of asthma medications at school and also to authorize sharing student health information with school staff.
- Provide this copy of the form to the school/childcare center/work/caretaker or other involved third party. (This copy may also be faxed to the school, etc.)

FOR MORE INFORMATION:

To access the August 2007 full version of the NHLBI Guidelines for the Diagnosis and Treatment of Asthma (EPR-3) or the October 2007 Summary Report, visit <http://www.nhlbi.nih.gov/guidelines/asthma/index.htm>.

THIS PAGE HAS BEEN INTENTIONALLY LEFT BLANK

My Asthma Plan

ENGLISH

Patient Name: _____

Medical Record #: _____

Provider's Name: _____

DOB: _____

Provider's Phone #: _____ Completed by: _____ Date: _____

Controller Medicines	How Much to Take	How Often	Other Instructions
		_____ times per day EVERY DAY!	<input type="checkbox"/> Gargle or rinse mouth after use
		_____ times per day EVERY DAY!	
		_____ times per day EVERY DAY!	
		_____ times per day EVERY DAY!	
Quick-Relief Medicines	How Much to Take	How Often	Other Instructions
<input type="checkbox"/> Albuterol (ProAir, Ventolin, Proventil) <input type="checkbox"/> Levalbuterol (Xopenex)	<input type="checkbox"/> 2 puffs <input type="checkbox"/> 4 puffs <input type="checkbox"/> 1 nebulizer treatment	Take ONLY as needed (see below — starting in Yellow Zone or before exercise)	NOTE: If you need this medicine more than two days a week, call physician to consider increasing controller medications and discuss your treatment plan.


Special instructions when I am



doing well,



getting worse,



having a medical alert.

GREEN ZONE

Doing *well*.

- No cough, wheeze, chest tightness, or shortness of breath during the day or night.
- Can do usual activities.



Peak Flow (for ages 5 and up):

is _____ or more. (80% or more of personal best)

Personal Best Peak Flow (for ages 5 and up): _____

PREVENT asthma symptoms every day:

- Take my controller medicines (above) every day.
- Before exercise, take _____ puff(s) of _____
- Avoid things that make my asthma worse.
(See back of form.)

YELLOW ZONE

Getting *worse*.



- Cough, wheeze, chest tightness, shortness of breath, or
- Waking at night due to asthma symptoms, or
- Can do some, but not all, usual activities.

Peak Flow (for ages 5 and up):

_____ to _____ (50 to 79% of personal best)

CAUTION. Continue taking every day controller medicines, AND:

- Take _____ puffs or _____ one nebulizer treatment of quick relief medicine. If I am not back in the **Green Zone** within 20-30 minutes take _____ more puffs or nebulizer treatments. If I am not back in the **Green Zone** within one hour, then I should:
- Increase _____
- Add _____
- Call _____
- Continue using quick relief medicine every 4 hours as needed. Call provider if not improving in _____ days.

RED ZONE

Medical Alert



- Very short of breath, or
- Quick-relief medicines have not helped, or
- Cannot do usual activities, or
- Symptoms are same or get worse after 24 hours in Yellow Zone.

Peak Flow (for ages 5 and up):

less than _____ (50% of personal best)

MEDICAL ALERT! Get help!

- Take quick relief medicine: _____ puffs every _____ minutes and get help immediately.
- Take _____
- Call _____

Danger! Get help immediately! Call 911 if trouble walking or talking due to shortness of breath or if lips or fingernails are gray or blue. For child, call 911 if skin is sucked in around neck and ribs during breaths or child doesn't respond normally.

Health Care Provider: My signature provides authorization for the above written orders. I understand that all procedures will be implemented in accordance with state laws and regulations. Student may self carry asthma medications: Yes No self administer asthma medications: Yes No (This authorization is for a maximum of one year from signature date.)

Healthcare Provider Signature

Date

ORIGINAL (Patient) / CANARY (School/Child Care/Work/Other Support Systems) / PINK (Chart)

This Asthma Plan was developed by a committee facilitated by the Regional Asthma Management and Prevention (RAMP) Initiative, a program of the Public Health Institute. This publication was supported by Cooperative Agreement Number 1U58DP001016-01 from the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of CDC. This plan is based on the recommendations from the National Heart, Lung, and Blood Institute's, "Guidelines for the Diagnosis and Management of Asthma," NIH Publication No. 07-4051 (August 2007). The information contained herein is intended for the use and convenience of physicians and other medical personnel and may not be appropriate for use in all circumstances. Decisions to adopt any particular recommendation must be made by qualified medical personnel in light of available resources and the circumstances presented by individual patients. No entity or individual involved in the funding or development of this plan makes any warranty or guarantee, express or implied, of the quality, fitness, performance or results of use of the information or products described in the plan or the Guidelines.

For additional information, please contact
RAMP at (510) 302-3365, <http://www.rampasthma.org>.

Cách Thức Phòng Ngừa Bệnh Suyễn

VIETNAMESE

Patient Name: _____

Medical Record #: _____

Tên của Nhà cung cấp dịch vụ chăm sóc sức khỏe: _____ DOB: _____

Số điện thoại của Nhà cung cấp dịch vụ chăm sóc sức khỏe: _____ Người điền đơn: _____ Ngày: _____

Các thuốc suyễn phải dùng mỗi ngày	Liều dùng	Số lần dùng	Các hướng dẫn Khác
		Lần/ngày MỖI NGÀY!	<input type="checkbox"/> Súc hoặc rửa miệng sau khi sử dụng
		Lần/ngày MỖI NGÀY!	
		Lần/ngày MỖI NGÀY!	
		Lần/ngày MỖI NGÀY!	
Các thuốc Làm giảm Cơn suyễn Nhanh	Liều dùng	Số lần dùng	Các hướng dẫn Khác
<input type="checkbox"/> Albuterol (ProAir, Ventolin, Proventil) <input type="checkbox"/> Levalbuterol (Xopenex)	<input type="checkbox"/> 2 lần xịt <input type="checkbox"/> 4 lần xịt <input type="checkbox"/> 1 lần điều trị xông thuốc	CHỈ dùng khi cần (xem dưới đây — bắt đầu ở vùng màu vàng hoặc trước khi vận động)	GHI CHÚ: Nếu quý vị cần thuốc này cho hơn hai ngày một tuần, hãy gọi bác sĩ để xem xét tăng liều thuốc kiểm soát bệnh suyễn và thảo luận về kế hoạch điều trị của quý vị.

Hướng dẫn đặc biệt



Lúc sức khỏe tốt



Khi sức khỏe xấu đi



Khi có triệu chứng báo động

VÙNG MÀU XANH

Khi khỏe.

- Không ho, thở khò khè, tức ngực, thở gấp ban ngày hoặc ban đêm.
- Có thể làm những việc thông thường

Thông lượng Định (cho trẻ 5 tuổi trở lên):
là _____ hoặc lớn hơn. (80% hoặc lớn hơn thông lượng định tốt nhất cho cá nhân)

Thông lượng định tốt nhất cho cá nhân (cho trẻ 5 tuổi trở lên):



PHÒNG NGỪA những triệu chứng suyễn hàng ngày:

- Dùng thuốc điều trị bệnh suyễn của bạn.
- Trước khi tập thể dục, xịt thuốc _____ lần _____
- Tránh những điều làm bệnh suyễn nặng thêm.
(Xem mặt sau của mẫu này.)

VÙNG MÀU VÀNG

Khi sức khỏe xấu đi



- Ho, thở khò khè, tức ngực, thở gấp hoặc
- Thức giấc ban đêm do những triệu chứng suyễn, hoặc
- Có thể làm một số, nhưng không phải tất cả, những việc thông thường

Thông lượng định (cho trẻ 5 tuổi trở lên):
_____ đến _____ (50 đến 79% thông lượng định tốt nhất cho cá nhân)

CHÚ Ý Tiếp tục dùng thuốc kiểm soát bệnh suyễn mỗi ngày, VÀ:

- Xịt _____ lần hoặc _____ lần điều trị xông thuốc làm giảm cơn suyễn nhanh chóng. Nếu tôi không trở lại **vùng màu xanh** trong vòng 20-30 phút sau khi _____ lần xịt thêm hoặc điều trị xông thuốc. Nếu không trở lại **vùng màu xanh** trong vòng một giờ, thì tôi nên:
 - Tăng liều dùng _____
 - Thêm liều dùng _____
 - Liên lạc _____
 - Tiếp tục dùng thuốc giảm cơn suyễn nhanh cứ mỗi 4 giờ một lần theo nhu cầu. Hãy gọi nhà cung cấp dịch vụ chăm sóc sức khỏe không thuyên giảm trong _____ ngày.

VÙNG MÀU ĐỎ

Triệu chứng báo động



- Thở rất gấp, hoặc
 - Thuốc giảm cơn suyễn nhanh không có tác dụng, hoặc
 - Không thể làm những việc thông thường, hoặc
 - Các triệu chứng không thay đổi hoặc xấu hơn sau 24 giờ trong Vùng Màu Vàng.
- Thông lượng định** (cho trẻ 5 tuổi trở lên):
Thấp hơn _____ (50% thông lượng định tốt nhất cho cá nhân)

TRIỆU CHỨNG BÁO ĐỘNG! Liên lạc giúp đỡ!

- Dùng thuốc giảm cơn suyễn nhanh: _____ mỗi lần _____ phút và cần được giúp đỡ ngay.
- Dùng _____
- Gọi _____

Khẩn Cấp! Cần được giúp đỡ ngay! Hãy gọi 911 nếu đã ho hoặc nói khó khăn do thở gấp hoặc nếu môi hoặc ngón tay bị xám hoặc xanh. Đối với trẻ em, gọi 911 nếu da bị hút vào quanh cổ và sườn trong khi thở hoặc trẻ em không đáp ứng bình thường.

Health Care Provider: My signature provides authorization for the above written orders. I understand that all procedures will be implemented in accordance with state laws and regulations. Student may self carry asthma medications: Yes No self administer asthma medications: Yes No (This authorization is for a maximum of one year from signature date.)

Coi Chừng Những Nguyên Nhân Làm Bệnh Suyễn Trở Nặng

HÚT THUỐC LÁ

- Không hút thuốc lá. Tha gia các lớp giúp cai nghiện thuốc lá.
- Không cho phép hút thuốc lá trong nhà hoặc xe hơi mùi khói thuốc còn lại có thể gây ra cơn suyễn.
- Tránh xa người đang hút thuốc lá.
- Nếu quý vị hút thuốc lá, hãy hút ở bên ngoài.



BỤI

- Hút bụi hàng tuần bằng máy hút bụi có lọc bụi nồng suất cao hoặc máy hút bụi trung tâm. Cố gắng hút bụi trong khi người bệnh suyễn không có mặt trong nhà.
- Thay thảm, nếu được. Làm ẩm thảm trước khi thay và lau khô nền hoàn toàn.
- Lau nền bằng vải ẩm hàng tuần.
- Giặt giường và đồ chơi nhồi bông bằng nước nóng mỗi 1-2 tuần. Làm lạnh đồ chơi nhồi bông không giặt được trong 24 giờ.
- Bọc nệm và gối trong các bao chống bụi.
- Giảm ồn ào và loại bỏ đồ chơi nhồi bông, nhất là quanh giường.
- Thường xuyên thay các bộ lọc của hệ thống sưởi.



SÂU BỌ

- Không để thực phẩm hoặc rác thải ra ngoài. Bỏ thực phẩm trong hộp có nắp đóng chặt.
- Dùng bẫy và mồi tẩm thuốc độc, như acid boric để diệt dán. Thay vì dùng thuốc phun/diệt, đặt bẫy tránh xa trẻ em, như sau tủ lạnh.
- Hút xác dán và trám lỗ bằng chất trám hoặc lớp luaré đồng.
- Sửa ống nước rò rỉ, mái nhà dột, và các nguồn nước khác.



NẤM MÓC

- Sử dụng quạt hút hoặc mở cửa sổ cho thông gió theo chiều ngang khi tắm vòi hoa sen hoặc nấu nướng.
- Chùi sạch nấm móc trên những bề mặt cứng bằng bột giặt với nước nóng và chà bằng bàn chải cứng hoặc miếng bọt biển, rồi rửa sạch bằng nước. Vật liệu hấp thụ có nấm móc cần phải được thay thế.
- Bảo đảm không có người bệnh suyễn trong phòng khi chùi rửa.
- Sửa ống nước hoặc các nguồn nước khác hoặc nguồn hơi ẩm.



THÚ VẬT

- Cân nhắc không nuôi thú. Tránh thú có lông hoặc lông vũ.
- Giữ thú nuôi ngoài phòng ngủ của người bệnh suyễn.
- Rửa tay của quý vị và tay của người bệnh suyễn sau khi sờ thú.



MÙI/THUỐC XỊT

- Tránh sử dụng sản phẩm có mùi mạnh, như chất khử mùi và tạo mùi thơm trong nhà, và chất giặt rửa và các chất dùng cho vệ sinh cá nhân có mùi thơm.
- Không sử dụng nồi/lò hở để sưởi.
- Khi lau chùi, tránh xa người bệnh suyễn và không sử dụng các sản phẩm lau chùi có mùi mạnh.
- Tránh dùng sản phẩm phun mù.
- Tránh dùng sản phẩm lau chùi mạnh hoặc cực mạnh.
- Tránh dùng amoniac, thuốc tẩy và chất tịt trùng.



PHẦN HOA VÀ NẤM MÓC NGOÀI TRỜI

- Ở trong nhà khi có nhiều mốc và phấn hoa.
- Đóng cửa sổ trong mùa phấn hoa.
- Tránh dùng quạt; hãy dùng máy điều hòa không khí.

CẨM/CÚM

- Giữ cơ thể khỏe mạnh bằng cách tập thể dục và ngủ đầy đủ.
- Tránh tiếp xúc gần với người bị cảm cúm.
- Rửa tay thường xuyên và tránh sờ tay lên mặt.
- Chống ngừa cúm hàng năm.

THỜI TIẾT VÀ Ô NHIỄM KHÔNG KHÍ

- Nếu gặp vấn đề với khí lạnh, cố gắng thở bằng mũi thay vì bằng miệng và che khăn choàng.
- Tránh hóng gió ban ngày và ban đêm và tránh tập thể dục nặng vào những ngày đó.
- Vào những ngày thời tiết bị ô nhiễm nặng, hãy ở trong nhà đóng cửa sổ.

TẬP THỂ DỤC

- Khởi động trước khi tập thể dục.
- Phương pháp phòng ngừa bệnh suyễn hoạt động trong nhà thay thế vào những ngày nhiều phấn hoa hoặc ô nhiễm.
- Nếu được bác sĩ hướng dẫn, hãy dùng thuốc trước khi tập thể dục. (Xem vùng màu xanh của kế hoạch hành động đối với bệnh suyễn.)

My Asthma Plan

ENGLISH

Patient Name: _____

Medical Record #: _____

Provider's Name: _____

DOB: _____

Provider's Phone #: _____

Completed by: _____ Date: _____

Controller Medicines	How Much to Take	How Often	Other Instructions
		_____ times per day EVERY DAY!	<input type="checkbox"/> Gargle or rinse mouth after use
		_____ times per day EVERY DAY!	
		_____ times per day EVERY DAY!	
		_____ times per day EVERY DAY!	
Quick-Relief Medicines	How Much to Take	How Often	Other Instructions
<input type="checkbox"/> Albuterol (ProAir, Ventolin, Proventil) <input type="checkbox"/> Levalbuterol (Xopenex)	<input type="checkbox"/> 2 puffs <input type="checkbox"/> 4 puffs <input type="checkbox"/> 1 nebulizer treatment	Take ONLY as needed (see below — starting in Yellow Zone or before exercise)	NOTE: If you need this medicine more than two days a week, call physician to consider increasing controller medications and discuss your treatment plan.


Special instructions when I am



doing well,



getting worse,



GREEN ZONE

Doing *well*.

- No cough, wheeze, chest tightness, or shortness of breath during the day or night.
- Can do usual activities.



Peak Flow (for ages 5 and up):

is _____ or more. (80% or more of personal best)

Personal Best Peak Flow (for ages 5 and up): _____

PREVENT asthma symptoms every day:

- Take my controller medicines (above) every day.
- Before exercise, take _____ puff(s) of _____
- Avoid things that make my asthma worse.
(See back of form.)

YELLOW ZONE

Getting *worse*.



- Cough, wheeze, chest tightness, shortness of breath, or
- Waking at night due to asthma symptoms, or
- Can do some, but not all, usual activities.

Peak Flow (for ages 5 and up):

_____ to _____ (50 to 79% of personal best)

CAUTION. Continue taking every day controller medicines, AND:

- Take _____ puffs or _____ one nebulizer treatment of quick relief medicine. If I am not back in the **Green Zone** within 20-30 minutes take _____ more puffs or nebulizer treatments. If I am not back in the **Green Zone** within one hour, then I should:
- Increase _____
- Add _____
- Call _____
- Continue using quick relief medicine every 4 hours as needed. Call provider if not improving in _____ days.

RED ZONE

Medical Alert



- Very short of breath, or
- Quick-relief medicines have not helped, or
- Cannot do usual activities, or
- Symptoms are same or get worse after 24 hours in Yellow Zone.

Peak Flow (for ages 5 and up):

less than _____ (50% of personal best)

MEDICAL ALERT! Get help!

- Take quick relief medicine: _____ puffs every _____ minutes and get help immediately.
- Take _____
- Call _____

Danger! Get help immediately! Call 911 if trouble walking or talking due to shortness of breath or if lips or fingernails are gray or blue. For child, call 911 if skin is sucked in around neck and ribs during breaths or child doesn't respond normally.

Health Care Provider: My signature provides authorization for the above written orders. I understand that all procedures will be implemented in accordance with state laws and regulations. Student may self carry asthma medications: Yes No self administer asthma medications: Yes No (This authorization is for a maximum of one year from signature date.)

Healthcare Provider Signature

Date

ORIGINAL (Patient) / CANARY (School/Child Care/Work/Other Support Systems) / PINK (Chart)

MẪU ĐƠN ỦY QUYỀN CHO NHÀ TRƯỜNG — Do phụ huynh/ người giám hộ điền và gửi lại cho nhà trường

VIETNAMESE

GIÁY ỦY QUYỀN VÀ MIỄN TRÁCH NHIỆM CỦA PHỤ HUYNH/NGƯỜI GIÁM HỘ: Tôi yêu cầu nhà trường hỗ trợ con tôi dùng các thuốc trị bệnh suyễn được liệt kê trong mẫu đơn này, và kế hoạch hành động đối với Bệnh suyễn, theo theo luật và quy định của tiểu bang.

Có Không

Con tôi có thể mang theo và tự dùng thuốc suyễn, tôi đồng ý miễn cho nhân viên của học khu và nhà trường không bị ràng buộc trách nhiệm nếu con tôi bị ảnh hưởng không tốt của việc tự sử dụng thuốc trị suyễn:

Có Không

Chữ ký của phụ huynh/người giám hộ

Ngày

GIÁY ỦY QUYỀN SỬ DỤNG VÀ TIẾT LỘ CHI TIẾT SỨC KHỎE CHO HỌC KHU

Văn bản đầy đủ này cho phép tiết lộ và/hoặc sử dụng chi tiết sức khỏe có thể nhận diện cá nhân, như ghi rõ dưới đây, phù hợp với luật liên bang (kể cả HIPAA) về tính riêng tư của chi tiết đó. Việc không cung cấp tất cả thông tin theo yêu cầu có thể làm mất hiệu lực của giấy ủy quyền này.

SỬ DỤNG VÀ TIẾT LỘ CHI TIẾT SỨC KHỎE:

Tên của bệnh nhân/Học sinh _____ / _____
Họ _____ Tên _____ Tên lót _____ Ngày Sinh _____

Tôi, người ký tên dưới đây, bằng phương tiện này cho phép (tên của cơ quan và/hoặc nhà cung cấp dịch vụ chăm sóc sức khỏe):

(1) _____ (2) _____ để cung cấp chi tiết sức khỏe từ hồ sơ y tế của con tôi có tên ở trên cho và từ:

Trường và học khu được ủy quyền tiết lộ thông tin _____ Địa chỉ / Thành phố và Tiểu bang / Mã Zip _____

Người liên lạc của trường hoặc học khu _____ Mã khu vực và Số điện thoại _____

Việc tiết lộ chi tiết sức khỏe cần thiết cho những mục đích sau:

Chi tiết yêu cầu được giới hạn như sau: Tất cả chi tiết sức khỏe; hoặc Chi tiết cụ thể về căn bệnh như được mô tả:

THỜI HẠN:

Giấy ủy quyền này sẽ có hiệu lực tức thời và có hiệu lực đến _____ (nhập ngày) hoặc 1 năm kể từ ngày ký tên, nếu không nhập ngày.

GIỚI HẠN

Luật pháp nghiêm cấm người yêu cầu tiết lộ thêm chi tiết sức khỏe của bạn trừ khi người yêu cầu có thêm một giấy ủy quyền khác của bạn hoặc trừ khi sự tiết lộ được luật pháp cho phép.

QUYỀN CỦA QUÝ VỊ:

Tôi hiểu rằng những quyền sau đây liên quan đến giấy ủy quyền này: Tôi có thể thu hồi giấy ủy quyền này bất cứ lúc nào. Việc hủy bỏ ủy quyền của tôi phải bằng văn bản, được tôi ký tên hoặc người đại diện của tôi ký tên, và giao cho cơ quan/nhân viên chăm sóc sức khỏe được liệt kê ở trên. Việc hủy bỏ ủy quyền của tôi có hiệu lực khi người yêu cầu nhận được, nhưng sẽ không có hiệu lực khi người yêu cầu hoặc những người khác đã hành động theo giấy ủy quyền này.

TIẾT LỘ LẠI:

Tôi hiểu rằng người yêu cầu (học khu) sẽ bảo vệ thông tin này theo quy định của luật bảo vệ quyền bình đẳng trong gia đình (FERPA) và thông tin này trở thành một phần trong hồ sơ giáo dục của học sinh. Chi tiết này sẽ được chia sẻ bởi những người làm việc tại hoặc với học khu cho mục đích tạo ra môi trường giáo dục an toàn, thích hợp và hạn chế tối thiểu, và các dịch vụ và chương trình y tế học đường.

Tôi có quyền nhận một bản sao giấy ủy quyền này. Có thể cần phải ký tên vào giấy ủy quyền này để học sinh nhận được những dịch vụ thích hợp trong môi trường giáo dục.

CHẤP THUẬN: _____
Tên viết bằng chữ in _____ Chữ ký _____ Ngày _____

Quan hệ với Bệnh nhân/học sinh

Mã khu vực và Số điện thoại